



Sacred Heart
UNIVERSITY

DEPARTMENT OF BIOLOGY

February 20, 2022

To: Commissioner Dykes, CT-DEEP

From: Jennifer H. Mattei, Ph.D. Professor, Department of Biology, IUCN Species Survival Commission Horseshoe Crab Specialist Group, Steering Committee Member

Regarding: Response to call from CTDEEP on public hearing regarding H.B. 5140 AN ACT CONCERNING THE HAND-HARVESTING OF HORSESHOE CRABS IN THE STATE

Horseshoe Crab harvest regulations.

Executive Summary of 20 years of research on the Population Ecology of the American Horseshoe Crab population inhabiting Long Island Sound.

I have been conducting research on the population ecology of *Limulus polyphemus*, the American horseshoe crab, for over 20 years. In short, due to multiple stressors, the population is in decline and due to very low population numbers the horseshoe crab is functionally extinct in Long Island Sound.

I first brought this to the attention of CTDEEP in 2005 when it was decided that three areas would be off limits to harvest. Our research has established long-term census records of hundreds of spawning horseshoe crabs at two of the protected sites, i.e. Sandy Point in West Haven and Milford Point in Milford. The third no harvest area, Menunketesuck Island, has not been surveyed. Since it takes 10 to 12 years for juvenile horseshoe crabs to develop into adults, it would be difficult to assess if this conservation effort had any measurable effect on the population until now. Since 2006, we have continued to monitor the spawning horseshoe crabs in these 2 areas and across beaches in Connecticut with our well-known citizen science program, Project *Limulus*. We have found that these two no harvest zones have **not** resulted in an increase in the spawning horseshoe crab population. In fact, we have documented the continued decline of this species (see Fig. 1).

Even though only ~12 permits exist to harvest horseshoe crabs and the reported catch is low, thousands of additional crabs are harvested every spawning season (Mattei, personal observation and observations reported by citizen scientists of Project *Limulus*). The hand harvest of the breeding population is a major cause of population decline in the Sound. Other stressors include loss of habitat (both marsh and sandy areas), pollution, bycatch in ghost nets and abandoned lobster traps as well as entrapment in intake pipes of powerplants (Smith et al 2017, Mattei, pers. obs.).

In 2015, we recommended changing the harvest regulations to a male only harvest which is the practice in Delaware (Beekey and Mattei, 2015). Now, we highly recommend closing all harvest of this species in Connecticut and New York (Mattei et al. in press). Our tag data has shown that across the beaches in Connecticut, the number of mated pairs found is declining and the number of single females found on the beach without a mate is increasing. The density of spawning horseshoe crabs is so low that the females cannot find mates and therefore this population is not reproducing at its maximum potential. The horseshoe crabs in Long Island Sound are not endangered of extirpation, at least currently, but their ecological role has been severely diminished. Horseshoe crabs function as a dominant species (not a keystone species) and their abundance is what is of ecological importance. They are no longer a source of food for shorebirds and fish

in LIS. No eggs are washed into the surf. Yes, you do see thousands of horseshoe crabs in some areas of Connecticut but their population density does not allow them to function within the food web as they have in the past. This “functional extinction” has occurred with other species in CT as well including pollinating insects, oysters, *Spartina* grasses, etc. **A management plan that merely keeps a species from going extinct is not good enough. It should keep the Long Island Sound ecosystem functioning sustainably.**




Figure 1. Average nighttime spawning index/m² ±SE for all surveyed beaches from 2008-2018. Parentheses indicate the number of surveys included in the average. $Y = -0.0013x + 0.0291$, $R^2 = 0.2609$, $p > 0.05$ (Mattei et al. in press).

References.

- Mattei, J.H., J.M. Kasinak, S.A. Senbel and K. Bartholomew. (In Press; 2022). The Power of Citizen Science: twenty years of horseshoe crab community research merging conservation, education and management. In Tanacredi, JT, Botton ML, Shin P, Mattei, JH, Cheung SG, Kwan KY, and Iwasaki Y. (eds.) *International Horseshoe Crab Conservation and Research Efforts: 2007- 2020*. Springer Pub. New York, NY.
- Youseff, I., S.A. Senbel, J. Kasinak, J. Mattei, 2019. RCrab: An R Analytics Tool to Visualize and Analyze the Movement of Horseshoe Crabs in Long Island Sound. Institute of Electrical and Electronics Engineers (IEEE) Long Island Systems, Applications, and Technology Conference (LISAT) Proceedings, May 3rd.
- Kasinak, J. J. Mattei 2017. Where have all the baby horseshoe crabs gone? Wrack Lines. Spring/Summer pg. 22-23, CT Sea Grant.
- Mattei, JH, M. Botton, M. Beekey, and C. Colón. 2015. Horseshoe Crab Research in Urban

- Estuaries: Challenges and Opportunities. In Carmichael, R, Botton ML, Cheung SG (eds.) *Changing global perspectives on biology, conservation, and management of horseshoe crabs*. Springer Pub. New York, NY.
- Beekey, MA, JH Mattei. 2015. The Mismanagement of *Limulus polyphemus* in Long Island Sound: What are the Characteristics of a Population in Decline? In Carmichael, R, Botton ML, Cheung SG (eds.) *Changing global perspectives on biology, conservation, and management of horseshoe crabs*. Springer Pub. New York, NY.
- Mattei, J. 2015. *Seas of Change: The Restoration of Ecosystem Services in Long Island Sound*. Wrack Lines. Fall/Winter pg. 22-23, CT Sea Grant.
<http://seagrant.uconn.edu/publications/magazines/wracklines/fallwinter15/index.php>
- Beekey, M.A., J.H. Mattei and B.J. Pierce. 2013. Horseshoe crab eggs: A rare resource for predators in Long Island Sound. *Journal of Experimental Marine Biology and Ecology* 439:152-159.
- Mattei J.H., M.A. Beekey, A. Woronik, H. Potter, C. Bond, K. Smith. 2011. Estimation of short-term tag mortality in the American horseshoe crab, *Limulus polyphemus*. *Transactions of the American Fisheries Society* 140:(4) 954-958.
- Kasinak, J., K. Bartholomew, M.A. Beekey and J.H. Mattei. 2011. Movement Patterns and Population Genetics of the American Horseshoe Crab in Relation to Long Island Sound Conservation Strategies. *Proceedings of the 2010 Biennial Long Island Sound Research Conference*. University of Connecticut, Stamford, CT.pg.20-25.
- Mattei, J.H., M.A. Beekey, A. Rudman, A. Woronik. 2010. Reproductive behavior in horseshoe crabs: does density matter? *Current Zoology* 56(5):634-642.
- Smith, D.R., H.J. Brockman, M.A. Beekey, T.L. King, M.J. Millard, J. Zaldívar-Rae. 2017. Conservation status of the American horseshoe crab, (*Limulus polyphemus*): a regional assessment. *Reviews in Fish Biology and Fisheries*, 27(1):135-1

Jennifer H. Mattei, Ph.D.
 Professor
 Department of Biology
 Sacred Heart University
 5151 Park Avenue
 Fairfield, CT 06825
 203-365-7577
matteij@sacredheart.edu
www.projectlimulus.org
www.sacredheart.edu/livingshorelines